



CENTER FOR DIGESTIVE HEALTH & NUTRITION  
725 CHERRINGTON PARKWAY • MOON TOWNSHIP, PA 15108

## COVID-19 2023 NOTICE

If you have a fever, unexplained cough, or respiratory symptoms or have been exposed to someone with symptoms or confirmed COVID-19 infection in the last 21 days, CALL and reschedule your colonoscopy procedure. Don't do the prep and risk being turned away.

## NuLYTELY (4 Liter PEG) Colonoscopy Prep

Modern colonoscopy is a diagnostic procedure to examine the colon or large intestine. Over the past 40 years, it has become the gold standard in screening for colon polyps and prevention of colorectal cancer.

**WARNING:** Individuals taking the semaglutide class of medications known as (GLP-1) receptor agonists may delay how their stomach empties. This may interfere with colonoscopy preparation AND increase the risk of aspiration under anesthesia for BOTH colonoscopy and upper endoscopy (EGD) exams.

ALL patients taking these medications once weekly MUST stop them temporarily for 1 week before their procedure appointment.

ALL patients taking these medications daily MUST stop them temporarily for 24 hours before their procedure appointment.

Failure to do so will result in the cancellation of the procedure, which will have to be rescheduled.



Most exams are quick, painless, and very safe with modern sedation methods and newer thin, flexible instruments. For an accurate and safe exam, however, it is very important that your colon is purged of all waste material by means of a restricted diet and laxative preparation, or "prep." Colonoscopy is an important test. Read this carefully and take the time to prepare correctly. REMEMBER: An accurate colonoscopy requires an excellent prep.

## **Special Instruction for Diabetics (READ LAST PAGE)**

We realize our patients with diabetes may be wondering about the safety of a colonoscopy prep. The bottom line is it can be done safely if you follow instructions. If you are taking medication for Diabetes, please read the last page for special instructions

### **THREE (3) DAYS BEFORE YOUR COLONOSCOPY**

Make arrangements for a companion to accompany you to your appointment and drive you home. It is best that this person is someone we can share the examination results with. *Bring a driver who can remain in the PARKING LOT (due to Covid) the entire time you are here. Arriving late or without a driver may require canceling and rescheduling your procedure.*

Eat normally, but don't overeat to compensate for the upcoming fast. Please avoid Metamucil, vitamins containing iron, raw vegetables, corn, and foods with small seeds such as tomatoes, sesame seeds, kiwi, and cucumbers for a few days before your exam. Go to the drug store and purchase your prep kit with the attached prescription if this was not already called into your pharmacy.

- one 4-liter bottle of NuLYTELY lavage solution ( or GoLytely , Colyte, PEG, etc)

**For this preparation, you may use any brand of PEG lavage solution.** None can claim to taste good, but probably the least offensive is NuLYTELY which is less salty and comes with flavor packs. Other commonly prescribed solutions include GoLytely, CoLyte, and PEG generic. They all work the same way. Some come pre-flavored. Golytely and generic PEG are unflavored, but you may add some lemon or iced tea-flavored Crystal Light.

Purchase the prep items at least one day before the day of preparation and read these instructions carefully before you begin.

### **TWO (2) DAYS BEFORE YOUR COLONOSCOPY**

*Regular activity and diet, but don't overeat. Also, don't go into the prep constipated.* If you feel that is the case, we suggest you take an over-the-counter laxative (such as 4 TBSP of Milk of Magnesia) and 4 glasses of water after dinner TWO nights before your exam to “prime the pump.”



We also recommend that you also purchase:

- A soft brand of toilet tissue. We suggest Charmin Plus.
- A box of baby wipes such as Huggies or Pampers Aloe wipes
- One small tube of diaper rash or Zinc Oxide cream (such as A & D Zinc Oxide)
- Two bottles of Boost Plus drink, (*Diabetics use Boost Glucose Control.*)
- A small box of Animal Crackers or Vanilla Wafers.
- Clear liquids and Jello. See list at end. (NO red, green, blue, or purple.)

### **ONE (1) DAY BEFORE YOUR COLONOSCOPY**

Food will be restricted this day, but you MUST drink plenty of clear fluids to prevent dehydration from the effects of the laxative. You may have as much Gatorade, clear broth, bouillon, iced tea, hot tea, black coffee, strained or clear fruit juices, flavored jello, sodas, Kool-Aid, clear hard candy, Italian ice, or popsicles as you like. Please avoid bright red, purple, or blue-colored items such as Cherry Jello or Cherry Popsicles. The best choice is banana, pineapple, orange, or lemon flavored. No solids. No milk or cream products are allowed.

Have a light breakfast such as one large or two medium eggs *or* a small portion of skinless chicken, turkey, or fish with 1 slice of white toast, pulp-free orange juice, and cream less coffee or tea. You may take all of your normal prescription medications. **If you are taking blood thinners such as Coumadin, Plavix, Ticlid, Aggrenox, or if you are taking Insulin or pills for diabetes, call for special instructions before you begin your prep.** Prepare the PEG solution according to the instructions on the bottle. (To improve the flavor of the unflavored lavage solution, you may add one packet of Lemonade or Ice Tea Flavored Crystal Light mix from your grocery.) Chill the solution in your refrigerator. You may go to work, but bring two bottles of Boost Plus and a box of animal crackers.

#### **Approximately 10 AM**

To minimize hunger and maintain your energy, have:

One 8 oz bottle of chilled Boost Plus or Ensure Plus and just four (4) Animal Crackers or Vanilla Wafers. Follow with 8 oz of any clear liquid.

#### **Approximately Noon**

To minimize hunger and maintain your energy, have another:

One 8 oz bottle of chilled Boost Plus or Ensure Plus and just four (4) Animal Crackers or Vanilla Wafers. Follow with 8 oz of any clear liquid.

#### **Approximately 6:00 PM**

Begin drinking the PEG lavage solution. You should drink 8 ounces every 15 minutes until half of the solution (8 glasses) is consumed. This will take about 2 hours. Drinking each glass quickly rather than slowly sipping it is best, although some patients prefer drinking it through a straw. It is better chilled. If you develop nausea or vomiting, pause for an hour and try to resume the preparation.

#### **Approximately 8:00 PM:**

You will have finished drinking half of the bottle. You can expect most of the fluid to be evacuated within an hour or so. You may drink as much clear liquid as you wish - the more, the better. Put the remaining PEG solution in your refrigerator overnight. Go to bed early and get a good night's sleep. Don't forget to set your alarm.

## **MORNING OF YOUR COLONOSCOPY**

### **Four and 1/2 Hours before your appointment**

Do NOT eat breakfast. Resume drinking the PEG lavage solution. Please drink four (4) more full glasses of the remaining solution. *You must finish this at least 4 hours before your exam and remain fasting the last 4 hours before your procedure time.* NO additional fluids. NO gum, candy, or mints. If you are diabetic, please do not take your diabetic medication on the morning of the exam. Unless otherwise instructed, you may take all of your other regular medications with a sip of water. **It is normal to continue to pass small amounts of clear or yellow liquid.**

You should plan to arrive at your scheduled appointment time. This allows time for registration, prep preparation, and any special orders your physician may require, and to answer any questions you may have.

Wear loose, comfortable, easily removable clothing and leave high heels, jewelry, and other valuables at home. **Please bring your insurance card and driver's license. If there is a deductible or copay, be prepared to pay it.**

Plan to arrive ONE HOUR before your procedure time.

Plan to arrive ON TIME for your appointment. Wear comfortable clothing and shoes that are easily removed. Leave jewelry and other valuables at home. No firearms are permitted. Bring your insurance card and driver's license. If there is a copay or deductible, be prepared to pay it PRIOR to your procedure.

## **YOUR DRIVER**

*Bring a driver who can remain here the entire time you are here.* The average visit is about two hours. Arriving late or without a driver may require canceling and rescheduling your procedure. Patients that would like to have their driver come to recovery after their procedure must have their driver remain in the waiting room the entire time. If the nurse comes to the waiting room to call your driver back and they have left the waiting room, they may miss the opportunity to come to recovery when the doctor goes over the results.

After the exam, you must be driven directly home, not to a restaurant. You may then gradually resume a normal diet and take a nap as the sedation wears off. You should not plan to work or drive until the following morning. You may resume normal activities the day after your exam. It is best that you do not plan to fly or travel long distances or to remote areas away from medical care for the first week, especially if a polyp was removed.

**Note: "CLEAR" LIQUIDS ALLOWED:** This includes clear fat-free broths, bouillon, coffee, and tea (without cream or non-dairy creamer), Kool-Aid, Crystal Light, carbonated beverages, clear sodas like ginger ale, orange juice, and other fruit juices without pulp, popsicles without pulp, Jell-O, hard candies, and Italian ices. **You are not allowed:** orange juice with pulp, fruit nectars, liquids you cannot see through, milk, cream, and non-dairy creamer. Avoid anything red or purple in color. No alcoholic drinks.

---

**YOU SHOULD CALL THE OFFICE AT 412-262-1000 AFTER 8:30 AM ON THE MORNING OF YOUR TEST IF YOU ARE STILL HAVING SOLID OR BROWN STOOLS.**

## **Three Rivers Endoscopy Center**

725 Cherrington Parkway • Moon Township, PA 15108-4305  
(412) 262-1000 IG: @thedigestivetract [www.gihealth.com](http://www.gihealth.com)

## Special Medication Instructions for Colonoscopy Exam

### Ozempic, Rybelsus, Wegovy, Mounjaro - Diabetic and Weight Loss Medications

IF daily capsule, stop 24 hours before the exam;

IF weekly injection, stop 1 week before the exam. Resume afterward at normal dose.

**WARNING:** Individuals taking the semiglutide class of medications known as (GLP-1) receptor agonists may delay how their stomach empties. This may interfere with colonoscopy preparation AND increase the risk of aspiration under anesthesia for BOTH colonoscopy and upper endoscopy (EGD) exams.

ALL patients taking these medications once weekly MUST stop them temporarily for 1 week before their procedure appointment.

ALL patients taking these medications daily MUST stop them temporarily for 24 hours before their procedure appointment.

Failure to do so will result in the cancellation of the procedure, which will have to be rescheduled.



### Metformin/Glucofage, Glucotrol/Glipizide, Trulicity, Byetta, Victoza, Januvia, Tradjenta

Stop the morning of the procedure and resume that evening.

### Jardiance, Forxiga, Brenzavvy, and Invokana

Stop taking three days before the procedure and resume once you're back to eating normally.

**Insulin** - Please speak to your endocrinologist or PCP, as the following are only starting points:

Regular insulin: People who use a "sliding scale" or insulin with meals can continue to follow their normal dosing strategy.

Long-acting insulin: Beginning the day before the procedure, patients will be asked to cut their basal insulin dose by 20 to 50 percent, depending on the insulin they use and the type of diabetes they experience.

### Anticoagulants "Blood thinners"

Coumadin, Warfarin, Jantofen, Eliquis, Xarelto, Pradaxa, Plavix, Aggrenox, Effient

Please speak to your physician or PCP because these drugs all act differently.



### Center For Digestive Health & Nutrition

725 Cherrington Parkway • Moon Township, PA 15108

412.262.1000 • [www.gihealth.com](http://www.gihealth.com) • IG: @thedigestivetract